

UBC ID #: \_\_\_\_\_ NAME (print): \_\_\_\_\_

Signature: \_\_\_\_\_



THE UNIVERSITY OF BRITISH COLUMBIA

IRVING K. BARBER SCHOOL  
OF ARTS AND SCIENCES  
UBC OKANAGAN

Instructor: Maria Martignoni Mseyá Course: MATH 225

Date: April 16th, 2019 Time: 1:00pm Duration: 3 hours

This exam has 10 questions for a total of 75 points.

### Rules governing a formal examination

- Each candidate must be prepared to produce a UBCcard for identification.
- Candidates are not permitted to ask questions of the invigilators, except in cases of supposed errors or ambiguities in examination questions.
- No candidate shall be permitted to enter the examination room after the expiration of one half-hour from the scheduled starting time, or to leave during the first half-hour of the examination.
- Candidates suspected of any of the following, or similar, dishonest practices shall be immediately dismissed from the examination and shall be liable to disciplinary action: having at the place of writing any books, papers or memoranda, calculators, computers, sound or image players/recorders/transmitters (including telephones), or other memory aid devices, other than those authorized by the examiners; speaking or communicating with other candidates; purposely exposing written papers to the view of other candidates or imaging devices.
- Candidates must not destroy or mutilate any examination material, must hand in all examination papers, and must not take any examination material from the examination room.

### Special Instructions

- Show and explain all of your work unless the question directs otherwise. Simplify all answers.
- Answer the questions in the spaces provided on the question sheets. If you run out of room for an answer, ask for extra paper.

This is a closed-book, individual exam.

Question:	1	2	3	4	5	6	7	8	9	10	Total
Points:	8	9	12	5	5	4	4	8	8	12	75
Score:											

- 8 1. Consider the equation

$$\frac{dv}{dt} = -a - kv$$

where  $a$  and  $k$  are positive constant parameters.

- (a) Find **all** solutions to the equation above.

(b) Find  $\lim_{t \rightarrow \infty} v(t)$

(c) Verify the answer with the use of a phase line diagram.

- 9 2. (a) Find the general solution to

$$\frac{d^2y}{dt^2} + 2\frac{dy}{dt} + 9y = 0 \quad (1)$$

- (b) Find the solution to Eq. (1) that satisfies the initial conditions  $y(0) = 1$ ,  $y'(0) = 0$ .

- (c) Sketch the solution you have obtained in part (b) as a function of time. Your sketch should include the initial conditions and possibly information about the location of maxima and/or minima.

- 12 3. (a) Find the general solution to

$$t^2 y'' - 4ty' + 6y = 0, \quad t > 0$$

- (b) Use the results obtained in part (a) to find the general solution to

$$t^2 y'' - 4ty' + 6y = t^3 + 1, \quad t > 0$$

Note: This is a NON-constant coefficient equation.

Hint: Remember to write the equation in standard form.

*Extra space for question 4*

- 5 4. Solve the following IVP by using the method of Laplace transform

$$y''' - y'' + y' - y = 0, \quad y(0) = 1, \quad y'(0) = 1, \quad y''(0) = 1$$

Hint:  $a^3 - a^2 + a - 1 = (a - 1)(a^2 + 1)$



5. (a) Write the following IVP

$$y' + \frac{3y}{x} + 2 = 3x \quad , \quad y(1) = 0$$

by using the Forward Euler formula with stepsize  $h$

- (b) Express  $y_1$  in terms of  $h$ . Simplify your answer and show all your work.

- 4 6. Verify that when the linear differential equation  $[P(x)y - Q(x)]dx + dy = 0$  is multiplied by  $\mu(x) = e^{\int P(x)dx}$ , the result is exact.

- 4 7. Use the convolution theorem to find the inverse Laplace transform of

$$F(s) = \frac{3s}{(s^2 + 1)(s + 4)}$$

(Do not solve the integral).

8. (a) Find the general solution to the differential equation  $y'' + 4y = 0$ . Express the general solution in the form  $y(t) = c_1y_1(t) + c_2y_2(t)$ .

(b) Use the Wronskian to show that  $y_1(t)$  and  $y_2(t)$  form a fundamental solutions set.

(c) Show that there is no solution that satisfies the conditions  $y(0) = 2$  and  $y(\pi/2) = 0$ .

(d) Explain briefly why your results in part (c) do not contradict the existence and uniqueness theorem (corresponding to Theorem 5, stated at the back of the exam).

9. The mixing tank in Figure 1 initially holds 500 L of water (there is initially no salt in the tank). For the first 30 min of operation, valve A is open, adding 5 L/min brine containing a 0.4 Kg/L salt concentration. After 30 min, valve A is closed and valve B is switched in, adding a 0.6 Kg/L concentration at 5 L/min. The exit valve C removes 5 L/min, thereby keeping the volume constant.




Figure 1: Mixing tank

- (a) Write down the IVP characterising the system (do NOT solve the problem).

- (b) Use the method of the Laplace transform to rewrite the IVP found in (a) into the s-space. Solve for  $X(s)$ .

12 10. (a) Solve the following IVP:

$$\frac{d^2y}{dt^2} + y = 5 \cos(t), \quad y(0) = 0, \quad y'(0) = 1 \quad (2)$$



*Extra space for question 10 (a)*

(b) Sketch the solution

(c) Explain briefly what is a possible physical interpretation of the solution you found in part (a). Refer to the forced mass-spring oscillator in your answer.

*Empty page*

## Additional Information

**Theorem 5** Suppose  $p(t)$ ,  $q(t)$ , and  $g(t)$  are continuous on an interval  $(a, b)$  that contains the point  $t_0$ . Then, for any choice of the initial values  $Y_0$  and  $Y_1$ , there exists a unique solution  $y(t)$  on the same interval  $(a, b)$  to the initial value problem

$$y''(t) + p(t)y'(t) + q(t)y(t) = g(t), \quad y(t_0) = Y_0, \quad y'(t_0) = Y_1.$$

### BRIEF TABLE OF LAPLACE TRANSFORMS

$f(t)$	$F(s) = \mathcal{L}\{f\}(s)$
1	$\frac{1}{s}, \quad s > 0$
$e^{at}$	$\frac{1}{s-a}, \quad s > a$
$t^n, \quad n = 1, 2, \dots$	$\frac{n!}{s^{n+1}}, \quad s > 0$
$\sin(bt)$	$\frac{b}{s^2 + b^2}, \quad s > 0$
$\cos(bt)$	$\frac{s}{s^2 + b^2}, \quad s > 0$
$e^{at}t^n, \quad n = 1, 2, \dots$	$\frac{n!}{(s-a)^{n+1}}, \quad s > a$
$e^{at} \sin(bt)$	$\frac{b}{(s-a)^2 + b^2}, \quad s > a$
$e^{at} \cos(bt)$	$\frac{s-a}{(s-a)^2 + b^2}, \quad s > a$
$u(t-a)$	$\frac{e^{-as}}{s}, \quad s > a$

### BRIEF TABLE OF PROPERTIES OF THE LAPLACE TRANSFORM

- P3:  $\mathcal{L}\{e^{at}f(t)\}(s) = \mathcal{L}\{f\}(s-a)$   
 P4:  $\mathcal{L}\{f'\}(s) = s\mathcal{L}\{f\}(s) - f(0)$   
 P6:  $\mathcal{L}\{f^{(n)}\}(s) = s^n\mathcal{L}\{f\}(s) - s^{n-1}f(0) - s^{n-2}f'(0) - \dots - f^{(n-1)}(0)$   
 P7:  $\mathcal{L}\{t^n f(t)\}(s) = (-1)^n \frac{d^n}{ds^n} (\mathcal{L}\{f\}(s))$

THEOREM: TRANSLATION IN  $t$ 

Let  $F(s) = \mathcal{L}\{f\}(s)$  exist for  $s > a \geq 0$ . If  $a$  is a positive constant, then

$$\mathcal{L}\{f(t-a)u(t-a)\}(s) = e^{-as}F(s),$$

and, conversely, an inverse Laplace transform of  $e^{-at}F(s)$  is given by

$$\mathcal{L}^{-1}\{e^{-as}F(s)\}(t) = f(t-a)u(t-a),$$

where  $u(t)$  is the unit step function.

## TRANSFORM OF A PERIODIC FUNCTION

If  $f$  has period  $T$  and is piecewise continuous on  $[0, T]$ , then the Laplace transforms

$$F(s) = \int_0^{\infty} e^{-st}f(t)dt \text{ and } F_T(s) = \int_0^T e^{-st}f(t)dt$$

are related by

$$F(s) = \frac{F_T(s)}{1 - e^{-sT}}.$$

## CONVOLUTION THEOREM

Let  $f(t)$  and  $g(t)$  be piecewise continuous on  $[0, \infty)$  and of exponential order  $\alpha$  and set  $F(s) = \mathcal{L}\{f\}(s)$  and  $G(s) = \mathcal{L}\{g\}(s)$ . Then

$$\mathcal{L}\{f * g\}(s) = F(s)G(s),$$

or, equivalently,

$$\mathcal{L}^{-1}\{F(s)G(s)\} = (f * g)(t).$$

## WINDOW FUNCTION

The window function  $\Pi_{a,b}$  can be defined in terms of step functions:

$$\Pi_{a,b}(t) = u(t-a) - u(t-b).$$