



a place of mind
THE UNIVERSITY OF BRITISH COLUMBIA

IRVING K. BARBER SCHOOL
OF ARTS AND SCIENCES
UBC OKANAGAN

Instructor: Rebecca Tyson Course: MATH 319
Date: Nov 6th, 2014 Time: 12:30pm Duration: 45 minutes.
This exam has 7 questions for a total of 35 points.

NAME: _____

SPECIAL INSTRUCTIONS

- Show and explain all of your work unless the question directs otherwise. Simplify all answers.
- The use of a calculator is permitted.
- Answer the questions in the spaces provided on the question sheets. If you run out of room for an answer, continue on the back of the page.

This is a two-stage exam. You have 45 minutes to complete the exam individually, then you will hand in the tests and join your group to redo the test as a group in the remaining 35 minutes.

Problem Number	1	2	3	4	5	6	Total
Points Earned							
Points Out Of	6	3	6	6	6	5	35

- 6 1. Consider the PDE problem below:

$$\begin{aligned}\frac{\partial u}{\partial t} &= \beta \frac{\partial^2 u}{\partial x^2}, & 0 < x < L, \quad t > 0, \\ u(0, t) &= U_1, \quad \frac{\partial u}{\partial x}(L, t) = U_2, & t > 0 \\ u(x, 0) &= f(x), & 0 < x < L.\end{aligned}$$

Note that U_1 and U_2 are constants. Derive and solve the steady state problem that arises from the PDE problem above.

- 3 2. Given that

$$\begin{aligned}\frac{\partial^2 u}{\partial t^2} &= 25 \frac{\partial^2 u}{\partial x^2}, & -\infty < x < \infty, \quad t > 0, \\ u_t(x, 0) &= 0, \quad u(x, 0) = f(x), & -\infty < x < \infty,\end{aligned}$$

what is the d'Alembert solution? Write it out algebraically, and then show what this solution means on the axis below by sketching the solution at $t=2$.




Figure 1: Plot of $f(x)$ for question 2.

- 6 3. Consider the BVP below:

$$F''(x) + \mu^2 F(x) = 0, \quad F'(0) = 0, \quad F(1) + F'(1) = 0.$$

Show that this BVP has multiple solutions (assume $\mu > 0$).

4. Laplace's equation on a wedge:

$$\frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} = 0, \quad 0 < r < 1, \quad 0 \leq \theta \leq \frac{\pi}{3}, \quad (1)$$

$$u(r, 0) = u\left(r, \frac{\pi}{3}\right) = 0, \quad u(1, \theta) = f(\theta), \quad 0 < r < 1, \quad 0 \leq \theta \leq \frac{\pi}{3}. \quad (2)$$

By assuming $u(r, \theta) = F(r)G(\theta)$, we reduce the PDE problem above to a pair of ODE problems. Solving, we find that the eigenvalues and eigenfunctions of the θ problem are

$$\lambda_n = (3n)^2, \quad G_n(\theta) = \sin(3n\theta), \quad n \in \mathbb{N} \quad (3)$$

The corresponding problem in r is

$$r^2 F'' + rF' - \lambda F = 0. \quad (4)$$

6 (a) Solve for $F(r)$.

3 (b) Write the formal solution for $u(r, \theta)$ and include the formula for the unknown Fourier coefficients.

- 6 5. Consider the functions

$$\left\{ \cos \left(\frac{n\pi x}{L} \right) \right\}_{n=1}^{\infty} \quad (5)$$

Show that these functions form an orthogonal set on the interval $0 \leq x \leq L$ with respect to the weight function $w(x) = 1$.

5 6. Given

$$u(x, t) = \sum_{n=1}^{\infty} c_n \sin\left(\frac{n\pi x}{L}\right) e^{-\alpha\left(\frac{n\pi}{L}\right)^2 t}, \text{ on } 0 < x < L, \quad (6)$$

where

$$u(x, 0) = f(x) = \frac{5}{3} \sin\left(\frac{6\pi x}{L}\right) + 3, \quad (7)$$

find the Fourier coefficients c_n and write out the full solution for $u(x, t)$.

7. (*Note: This problem is a BONUS on the GROUP test. If you have extra time, you can get started on it here.*)

Consider the following Dirichlet Laplace problem on a rectangular domain:

$$\nabla^2 u = 0, \quad 0 < x < a, \quad 0 < y < b, \quad (8)$$

$$u(0, y) = u(a, y) = 0, \quad 0 < y < b, \quad (9)$$

$$u(x, 0) = f_1(x), \quad u(x, b) = f_2(x), \quad 0 < x < a. \quad (10)$$

Find the formal solution for $u(x, y)$ (don't solve for the Fourier coefficients).

(Question 7 continued.)

Information That May Be Useful

Note: On this test, you may assume that for the BVP

$$F'' + \lambda F = 0,$$

the only nontrivial solutions are obtained for the case $\lambda = \mu^2 > 0$, and the corresponding solutions are

$$F(x) = c_1 \cos(\mu x) + c_2 \sin(\mu x),$$

without showing any work.

Possibly useful identities:

$$\cos(A) \cos(B) = \frac{1}{2} (\cos(A + B) + \cos(A - B))$$

$$\sin(A) \sin(B) = \frac{1}{2} (\cos(A - B) - \cos(A + B))$$

$$\sin(A) \cos(B) = \frac{1}{2} (\sin(A + B) + \sin(A - B))$$