

# Examples of discontinuous maximal monotone linear operators and the solution to a recent problem posed by B.F. Svaiter

Heinz H. Bauschke\*, Xianfu Wang† and Liangjin Yao‡

April 6, 2010

## Abstract

In this paper, we give two explicit examples of unbounded linear maximal monotone operators. The first unbounded linear maximal monotone operator  $S$  on  $\ell^2$  is skew. We show its domain is a proper subset of the domain of its adjoint  $S^*$ , and  $-S^*$  is not maximal monotone. This gives a negative answer to a recent question posed by Svaiter. The second unbounded linear maximal monotone operator is the inverse Volterra operator  $T$  on  $L^2[0, 1]$ . We compare the domain of  $T$  with the domain of its adjoint  $T^*$  and show that the skew part of  $T$  admits two distinct linear maximal monotone skew extensions. These unbounded linear maximal monotone operators show that the constraint qualification for the maximality of the sum of maximal monotone operators can not be significantly weakened, and they are simpler than the example given by Phelps-Simons. Interesting consequences on Fitzpatrick functions for sums of two maximal monotone operators are also given.

## 2000 Mathematics Subject Classification:

Primary 47A06, 47H05; Secondary 47A05, 47B65, 52A41.

**Keywords:** Adjoint operator, Fitzpatrick function, Fenchel conjugate, linear relation, maximal monotone operator, multifunction, monotone operator, skew operator, unbounded linear monotone operator.

---

\*Mathematics, Irving K. Barber School, UBC Okanagan, Kelowna, British Columbia V1V 1V7, Canada. E-mail: [heinz.bauschke@ubc.ca](mailto:heinz.bauschke@ubc.ca).

†Mathematics, Irving K. Barber School, UBC Okanagan, Kelowna, British Columbia V1V 1V7, Canada. E-mail: [shawn.wang@ubc.ca](mailto:shawn.wang@ubc.ca).

‡Mathematics, Irving K. Barber School, UBC Okanagan, Kelowna, British Columbia V1V 1V7, Canada. E-mail: [ljinyao@interchange.ubc.ca](mailto:ljinyao@interchange.ubc.ca).

# 1 Introduction

Linear monotone operators play important roles in modern monotone operator theory and partial differential equations [1, 2, 8, 14, 18, 22, 23, 24, 25], and they are examples that delineate the boundary of the general theory. In this paper, we explicitly construct two unbounded linear monotone operators (not full domain, linear and single-valued on their domains). They answer one of Svaiter's question, have some interesting consequences on Fitzpatrick functions for sums of two maximal monotone operators, and show that the constraint qualification for the maximality of the sum of maximal monotone operators can not be weakened significantly, see [16], [19, Theorem 5.5] and [22]. Our examples are simpler than the one given by [14].

The paper is organized as follows. Basic facts and auxiliary results are recorded in Section 2. In Section 3, we construct an unbounded maximal monotone skew operator  $S$  on  $\ell^2$ . For a maximal monotone skew operator, it is well known that its domain is always a subset of the domain of its adjoint. An interesting question remained is whether or not both of the domains are always same. The maximal monotone skew operator  $S$  enjoys the property that the domain of  $-S$  is a proper subset of the domain of its adjoint  $S^*$ , see Theorem 3.6. Svaiter asked in [21] whether or not  $-S^*$  (termed  $S^\dagger$  in [21]) is maximal monotone provided that  $S$  is maximal skew. This operator also answers Svaiter's question in the negative, see Theorem 3.15. In Section 4 we systematically study the inverse Volterra operator  $T$ . We show that  $T$  is neither skew nor symmetric and compare the domain of  $T$  with the domain of its adjoint  $T^*$ . It turns out that the skew part of  $T$ :  $S = \frac{T-T^*}{2}$  admits two distinct linear maximal monotone and skew extensions even the the domain of  $S$  is a dense linear subspace in  $L^2[0,1]$ . It was shown that Fitzpatrick functions  $F_{A+B} = F_A \square_2 F_B$  when  $A, B$  are maximal monotone linear relations and  $\text{dom } A - \text{dom } B$  is a closed subspace, see [5, Theorem 5.10]. Using these unbounded linear maximal monotone operators in Sections 3 and 4 we also show that the constraint qualification  $\text{dom } A - \text{dom } B$  being closed can not be significantly weakened either.

Throughout this paper, we assume that

$$X \text{ is a real Hilbert space, with inner product } \langle \cdot, \cdot \rangle.$$

Let  $S$  be a set-valued operator (also known as multifunction) from  $X$  to  $X$ . We say that  $S$  is *monotone* if

$$(\forall (x, x^*) \in \text{gra } S) (\forall (y, y^*) \in \text{gra } S) \quad \langle x - y, x^* - y^* \rangle \geq 0,$$

where  $\text{gra } S := \{(x, x^*) \in X \times X \mid x^* \in Sx\}$ ;  $S$  is said to be *maximal monotone* if no proper enlargement (in the sense of graph inclusion) of  $S$  is monotone. We say  $T$  is a maximal monotone extension of  $S$  if  $T$  is maximal monotone and  $\text{gra } T \supseteq \text{gra } S$ . The *domain* of  $S$  is  $\text{dom } S := \{x \in X \mid Sx \neq \emptyset\}$ , and its *range* is  $\text{ran } S := S(X) = \bigcup_{x \in X} Sx$ .

We say  $S$  is a *linear relation* if  $\text{gra } S$  is linear. The *adjoint* of  $S$ , written  $S^*$ , is defined by

$$\text{gra } S^* := \{(x, x^*) \in X \times X \mid (x^*, -x) \in (\text{gra } S)^\perp\},$$

where, for any subset  $C$  of a Hilbert space  $Z$ ,  $C^\perp := \{z \in Z \mid \langle z, c \rangle = 0, \forall c \in C\}$ . We say a linear relation  $S$  is *skew* if  $\langle x, x^* \rangle = 0, \forall (x, x^*) \in \text{gra } S$ , and  $S$  is a *maximal monotone skew operator* if  $S$

is a maximal monotone operator and  $S$  is skew. Svaiter introduced  $S^\dagger$  in [21], which is defined by

$$\text{gra } S^\dagger := \{(x, x^*) \in X \times X \mid (x^*, x) \in (\text{gra } S)^\perp\}.$$

Hence  $S^\dagger = -S^*$ . For each function  $f : X \rightarrow ]-\infty, +\infty]$ ,  $f^*$  stands for the *Fenchel conjugate* given by

$$f^*(x^*) = \sup_{x \in X} (\langle x^*, x \rangle - f(x)) \quad \forall x^* \in X.$$

## 2 Auxiliary results and facts

In this section we gather some facts about linear relations, monotone operators, and Fitzpatrick functions. They will be used frequently in sequel.

**Fact 2.1 (Cross)** *Let  $S : X \rightrightarrows X$  be a linear relation. Then the following hold.*

- (i)  $(S^*)^{-1} = (S^{-1})^*$ .
- (ii) *If  $\text{gra } S$  is closed, then  $S^{**} = S$ .*
- (iii) *If  $k \in \mathbb{R} \setminus \{0\}$ , then  $(kS)^* = kS^*$ .*
- (iv)  $(\forall x \in \text{dom } S^*)(\forall y \in \text{dom } S) \langle S^*x, y \rangle = \langle x, Sy \rangle$  *is a singleton.*

*Proof.* (i): See [10, Proposition III.1.3(b)]. (ii): See [10, Exercise VIII.1.12]. (iii): See [10, Proposition III.1.3(c)]. (iv): See [10, Proposition III.1.2]. ■

If  $S : X \rightrightarrows X$  is a linear relation that is at most single-valued, then we will identify  $S$  with the corresponding linear operator from  $\text{dom } S$  to  $X$  and (abusing notation slightly) also write  $S : \text{dom } S \rightarrow X$ . An analogous comment applies conversely to a linear single-valued operator  $S$  with domain  $\text{dom } S$ , which we will identify with the corresponding at most single-valued linear relation from  $X$  to  $X$ .

**Fact 2.2 (Phelps-Simons)** (See [14, Theorem 2.5 and Lemma 4.4].) *Let  $S : \text{dom } S \rightarrow X$  be monotone and linear. The following hold.*

- (i) *If  $S$  is maximal monotone, then  $\text{dom } S$  is dense (and hence  $S^*$  is at most single-valued).*
- (ii) *Assume that  $S$  is a skew operator such that  $\text{dom } S$  is dense. Then  $\text{dom } S \subseteq \text{dom } S^*$  and  $S^*|_{\text{dom } S} = -S$ .*

**Fact 2.3 (Brézis-Browder)** (See [9, Theorem 2].) *Let  $S : X \rightrightarrows X$  be a monotone linear relation such that  $\text{gra } S$  is closed. Then the following are equivalent.*

- (i)  $S$  is maximal monotone.
- (ii)  $S^*$  is maximal monotone.
- (iii)  $S^*$  is monotone.

For  $A: X \rightrightarrows X$ , the *Fitzpatrick function* associated with  $A$  is defined by

$$(1) \quad F_A: X \times X \rightarrow ]-\infty, +\infty]: (x, x^*) \mapsto \sup_{(a, a^*) \in \text{gra } A} (\langle x, a^* \rangle + \langle a, x^* \rangle - \langle a, a^* \rangle).$$

Following Penot [15], if  $F: X \times X \rightarrow ]-\infty, +\infty]$ , we set

$$(2) \quad F^\top: X \times X: (x^*, x) \mapsto F(x, x^*).$$

**Fact 2.4 (Fitzpatrick)** (See [12].) *Let  $A: X \rightrightarrows X$  be monotone. Then  $F_A = \langle \cdot, \cdot \rangle$  on  $\text{gra } A$  and  $F_{A^{-1}} = F_A^\top$ . If  $A$  is maximal monotone and  $(x, x^*) \in X \times X$ , then*

$$F_A(x, x^*) \geq \langle x^*, x \rangle,$$

*with equality if and only if  $(x, x^*) \in \text{gra } A$ .*

If  $A: X \rightarrow X$  is a linear operator, we write

$$(3) \quad A_+ = \frac{1}{2}A + \frac{1}{2}A^* \quad \text{and} \quad q_A: X \rightarrow \mathbb{R}: x \mapsto \frac{1}{2}\langle x, Ax \rangle.$$

**Fact 2.5** (See [4, Proposition 2.3] and [2, Proposition 2.2(v)]). *Let  $A: X \rightarrow X$  be linear and monotone, and let  $(x, x^*) \in X \times X$ . Then*

$$(4) \quad F_A(x, x^*) = 2q_{A_+}^*(\frac{1}{2}x^* + \frac{1}{2}A^*x) = \frac{1}{2}q_{A_+}^*(x^* + A^*x).$$

*If  $\text{ran } A_+$  is closed, then  $\text{dom } q_{A_+}^* = \text{ran } A_+$ .*

To study Fitzpatrick functions of sums of maximal monotone operator, one needs the  $\square_2$  operation:

**Definition 2.6** *Let  $F_1, F_2: X \times X \rightarrow ]-\infty, +\infty]$ . Then the partial inf-convolution  $F_1 \square_2 F_2$  is the function defined on  $X \times X$  by*

$$F_1 \square_2 F_2: (x, x^*) \mapsto \inf_{y^* \in X} (F_1(x, x^* - y^*) + F_2(x, y^*)).$$

**Fact 2.7** (See [18, Lemma 23.9] or [3, Proposition 4.2].) *Let  $A, B: X \rightrightarrows X$  be monotone such that  $\text{dom } A \cap \text{dom } B \neq \emptyset$ . Then  $F_A \square_2 F_B \geq F_{A+B}$ .*

Under some constraint qualifications, one has

**Fact 2.8** (i) (See [2].) *Let  $A, B: X \rightarrow X$  be continuous, linear, and monotone operators such that  $\text{ran}(A_+ + B_+)$  is closed. Then  $F_{A+B} = F_A \square_2 F_B$ .*

(ii) (See [5].) *Let  $A, B: X \rightrightarrows X$  be maximal monotone linear relations, and suppose that  $\text{dom } A - \text{dom } B$  is closed. Then  $F_{A+B} = F_A \square_2 F_B$ .*

### 3 An unbounded skew operator on $\ell^2$

In this section, we construct a maximal monotone and skew operator  $S$  on  $\ell^2$  such that  $-S^*$  is not maximal monotone. This answers one of Svaiter's question. We explicitly compute the Fitzpatrick functions  $F_{S+S^*}$ ,  $F_S$ ,  $F_{S^*}$ , and show that  $F_{S+S^*} \neq F_S \square_2 F_{S^*}$  even though  $S, S^*$  are linear maximal monotone with  $\text{dom } S - \text{dom } S^*$  being a dense linear subspace in  $\ell^2$ .

#### 3.1 The Example in $\ell^2$

Let  $\ell^2$  denote the Hilbert space of real square-summable sequences  $(x_n)_{n \in \mathbb{N}} = (x_1, x_2, x_3, \dots)$ , where  $\mathbb{N} = \{1, 2, 3, \dots\}$ .

**Example 3.1** Let  $X = \ell^2$ , and  $S : \text{dom } S \rightarrow \ell^2$  be given by

$$(5) \quad Sy := \frac{\left( \sum_{i < n} y_i - \sum_{i > n} y_i \right)_{n \in \mathbb{N}}}{2} = \left( \sum_{i < n} y_i + \frac{1}{2} y_n \right)_{n \in \mathbb{N}}, \quad \forall y = (y_n)_{n \in \mathbb{N}} \in \text{dom } S,$$

where  $\text{dom } S := \{y = (y_n)_{n \in \mathbb{N}} \in \ell^2 \mid \sum_{i \geq 1} y_i = 0, \left( \sum_{i \leq n} y_i \right)_{n \in \mathbb{N}} \in \ell^2\}$  and  $\sum_{i < 1} y_i = 0$ . In matrix form,

$$S = \frac{1}{2} \begin{pmatrix} 0 & -1 & -1 & -1 & -1 & \cdots & -1 & -1 & \cdots \\ 1 & 0 & -1 & -1 & -1 & \cdots & -1 & -1 & \cdots \\ 1 & 1 & 0 & -1 & -1 & \cdots & -1 & -1 & \cdots \\ 1 & 1 & 1 & 0 & -1 & \cdots & -1 & -1 & \cdots \\ 1 & 1 & 1 & 1 & 0 & \cdots & -1 & -1 & \cdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & & & & \ddots \end{pmatrix},$$

or

$$S = \begin{pmatrix} \frac{1}{2} & 0 & 0 & 0 & 0 & \cdots & 0 & 0 & \cdots \\ 1 & \frac{1}{2} & 0 & 0 & 0 & \cdots & 0 & 0 & \cdots \\ 1 & 1 & \frac{1}{2} & 0 & 0 & \cdots & 0 & 0 & \cdots \\ 1 & 1 & 1 & \frac{1}{2} & 0 & \cdots & 0 & 0 & \cdots \\ 1 & 1 & 1 & 1 & \frac{1}{2} & \cdots & 0 & 0 & \cdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & & & & \ddots \end{pmatrix}.$$

Using the second matrix, it is easy to see that  $S$  is injective.

**Proposition 3.2** *Let  $S$  be defined as in Example 3.1. Then  $S$  is skew.*

*Proof.* Let  $y = (y_n)_{n \in \mathbb{N}} \in \text{dom } S$ . Then  $\left( \sum_{i \leq n} y_i \right)_{n \in \mathbb{N}} \in \ell^2$ . Thus,

$$\ell^2 \ni \left( \sum_{i \leq n} y_i \right)_{n \in \mathbb{N}} - \frac{1}{2} y = \left( \sum_{i \leq n} y_i \right)_{n \in \mathbb{N}} - \frac{1}{2} (y_n)_{n \in \mathbb{N}} = \left( \sum_{i < n} y_i + \frac{1}{2} y_n \right)_{n \in \mathbb{N}} = Sy.$$

Hence  $S$  is well defined. Clearly,  $S$  is linear on  $\text{dom } S$ . Now we show  $S$  is skew.

Let  $y = (y_n)_{n \in \mathbb{N}} \in \text{dom } S$ , and  $s := \sum_{i \geq 1} y_i$ . Then  $\left( \sum_{i \leq n} y_i \right)_{n \in \mathbb{N}} \in \ell^2$ . Hence  $\left( \sum_{i < n} y_i \right)_{n \in \mathbb{N}} = \left( \sum_{i \leq n} y_i \right)_{n \in \mathbb{N}} - (y_n)_{n \in \mathbb{N}} \in \ell^2$ . By  $s = 0$ ,

$$\ell^2 \ni - \left( \sum_{i < n} y_i \right)_{n \in \mathbb{N}} = 0 - \left( \sum_{i < n} y_i \right)_{n \in \mathbb{N}} = \left( \sum_{i \geq 1} y_i - \sum_{i < n} y_i \right)_{n \in \mathbb{N}} = \left( \sum_{i \geq n} y_i \right)_{n \in \mathbb{N}},$$

$$(6) \quad \left( \sum_{i \geq n+1} y_i \right)_{n \in \mathbb{N}} = 0 - \left( \sum_{i \leq n} y_i \right)_{n \in \mathbb{N}} \in \ell^2.$$

Thus, by (6),

$$(7) \quad -2 \langle Sy, y \rangle = \left\langle \left( \sum_{i > n} y_i - \sum_{i < n} y_i \right)_{n \in \mathbb{N}}, y \right\rangle = \left\langle \left( \sum_{i \geq n+1} y_i + \sum_{i \geq n} y_i \right)_{n \in \mathbb{N}}, y \right\rangle$$

$$= \left\langle \left( \sum_{i \geq 1} y_i, \sum_{i \geq 2} y_i, \dots \right) + \left( \sum_{i \geq 2} y_i, \sum_{i \geq 3} y_i, \dots \right), y \right\rangle$$

$$= \langle (s, s - y_1, s - (y_1 + y_2), \dots) + (s - y_1, s - (y_1 + y_2), \dots), (y_1, y_2, \dots) \rangle$$

$$= [s y_1 + (s - y_1) y_2 + (s - (y_1 + y_2)) y_3 + \dots] +$$

$$[ (s - y_1) y_1 + (s - (y_1 + y_2)) y_2 + (s - (y_1 + y_2 + y_3)) y_3 + \dots ]$$

$$= \lim_n [s y_1 + (s - y_1) y_2 + \dots + (s - (y_1 + \dots + y_{n-1})) y_n] +$$

$$\lim_n [(s - y_1) y_1 + (s - (y_1 + y_2)) y_2 + \dots + (s - (y_1 + \dots + y_n)) y_n]$$

$$= \lim_n [s(y_1 + \dots + y_n) - y_1 y_2 - (y_1 + y_2) y_3 - \dots - (y_1 + \dots + y_{n-1}) y_n] +$$

$$[s(y_1 + \dots + y_n) - (y_1^2 + \dots + y_n^2) - y_1 y_2 - \dots - (y_1 + \dots + y_{n-1}) y_n]$$

$$= \lim_n [2s(y_1 + \dots + y_n) - (y_1 + \dots + y_n)^2] = 2s^2 - s^2 = s^2 = 0.$$

Hence  $S$  is skew. ■

**Remark 3.3**  $S$  is unbounded in Example 3.1, since  $(1, 0, 0, \dots, 0, \dots) \notin \text{dom } S$ .

**Fact 3.4 (Phelps-Simons)** (See [14, Proposition 3.2(a)]). *Let  $S : \text{dom } S \rightarrow X$  be linear and monotone. Then  $(x, x^*) \in X \times X$  is monotonically related to  $\text{gra } S$  if, and only if*

$$\langle x, x^* \rangle \geq 0 \text{ and } [\langle Sy, x \rangle + \langle x^*, y \rangle]^2 \leq 4 \langle x^*, x \rangle \langle Sy, y \rangle, \quad \forall y \in \text{dom } S.$$

**Proposition 3.5** *Let  $S$  be defined as in Example 3.1. Then  $S$  is a maximal monotone operator. In particular,  $\text{gra } S$  is closed.*

*Proof.* By Proposition 3.2,  $S$  is skew. Let  $(x, x^*) \in X \times X$  be monotonically related to  $\text{gra } S$ . Write  $x = (x_n)_{n \in \mathbb{N}}$  and  $x^* = (x_n^*)_{n \in \mathbb{N}}$ . By Fact 3.4, we have

$$(8) \quad \langle Sy, x \rangle + \langle x^*, y \rangle = 0, \quad \forall y \in \text{dom } S.$$

Let  $e_n = (0, \dots, 0, 1, 0, \dots)$ : the  $n$ th entry is 1 and the others are 0. Then let  $y = -e_1 + e_n$ . Thus  $y \in \text{dom } S$  and  $Sy = (-\frac{1}{2}, -1, \dots, -1, -\frac{1}{2}, 0, \dots)$ . Then by (8),

$$(9) \quad -x_1^* + x_n^* - \frac{1}{2}x_1 - \frac{1}{2}x_n - \sum_{i=2}^{n-1} x_i = 0 \Rightarrow x_n^* = x_1^* - \frac{1}{2}x_1 + \sum_{i=1}^{n-1} x_i + \frac{1}{2}x_n.$$

Since  $x^* \in \ell^2$  and  $x \in \ell^2$ , we have  $x_n^* \rightarrow 0, x_n \rightarrow 0$ . Thus by (9),

$$(10) \quad -\sum_{i \geq 1} x_i = x_1^* - \frac{1}{2}x_1.$$

Next we show  $-\sum_{i \geq 1} x_i = x_1^* - \frac{1}{2}x_1 = 0$ . Let  $s = \sum_{i \geq 1} x_i$ . Then by (9) and (10),

$$(11) \quad \begin{aligned} 2x^* &= 2(x_n^*)_{n \in \mathbb{N}} = 2 \left( -\sum_{i \geq 1} x_i + \sum_{i < n} x_i + \frac{1}{2}x_n \right)_{n \in \mathbb{N}} = \left( -2 \sum_{i \geq 1} x_i + 2 \sum_{i < n} x_i + x_n \right)_{n \in \mathbb{N}} \\ &= \left( -2 \sum_{i \geq n} x_i + x_n \right)_{n \in \mathbb{N}} = \left( -\sum_{i \geq n} x_i - \sum_{i \geq n} x_i + x_n \right)_{n \in \mathbb{N}} \\ &= \left( -\sum_{i \geq n} x_i - \sum_{i \geq n+1} x_i \right)_{n \in \mathbb{N}}. \end{aligned}$$

On the other hand, by (9),

$$\ell^2 \ni x^* - \frac{1}{2}x = \left( -\sum_{i \geq 1} x_i + \sum_{i < n} x_i + \frac{1}{2}x_n \right)_{n \in \mathbb{N}} - \left( \frac{1}{2}x_n \right)_{n \in \mathbb{N}} = \left( -\sum_{i \geq n} x_i \right)_{n \in \mathbb{N}}.$$

Then by (11),

$$2x^* = \left( -\sum_{i \geq n} x_i \right)_{n \in \mathbb{N}} + \left( -\sum_{i \geq n+1} x_i \right)_{n \in \mathbb{N}}.$$

Then by Fact 3.4, similar to the proof in (7) in Proposition 3.1, we have

$$\begin{aligned} 0 &\geq -2\langle x^*, x \rangle = \left\langle \left( \sum_{i \geq n} x_i \right)_{n \in \mathbb{N}} + \left( \sum_{i \geq n+1} x_i \right)_{n \in \mathbb{N}}, x \right\rangle \\ &= \left\langle \left( \sum_{i \geq 1} x_i, \sum_{i \geq 2} x_i, \dots \right) + \left( \sum_{i \geq 2} x_i, \sum_{i \geq 3} x_i, \dots \right), x \right\rangle \\ &= 2s^2 - s^2 = s^2. \end{aligned}$$

Hence  $s = 0$ , i.e.,  $x_1^* = \frac{1}{2}x_1$ . By (9),  $x^* = \left( \sum_{i < n} x_i + \frac{1}{2}x_n \right)_{n \in \mathbb{N}}$ . Thus

$$\ell^2 \ni x^* + \frac{1}{2}x = \left( \sum_{i < n} x_i + \frac{1}{2}x_n \right)_{n \in \mathbb{N}} + \left( \frac{1}{2}x_n \right)_{n \in \mathbb{N}} = \left( \sum_{i \leq n} x_i \right)_{n \in \mathbb{N}}.$$

Hence  $x \in \text{dom } S$  and  $x^* = Sx$ . Thus,  $S$  is maximal monotone. Hence  $\text{gra } S$  is closed. ■

**Proposition 3.6** *Let  $S$  be defined as in Example 3.1. Then*

$$(12) \quad S^*y = \left( \sum_{i>n} y_i + \frac{1}{2}y_n \right)_{n \in \mathbb{N}}, \quad \forall y = (y_n)_{n \in \mathbb{N}} \in \text{dom } S^*,$$

where  $\text{dom } S^* = \{y = (y_n)_{n \in \mathbb{N}} \in \ell^2 \mid \sum_{i \geq 1} y_i \in \mathbb{R}, \left( \sum_{i>n} y_i \right)_{n \in \mathbb{N}} \in \ell^2\}$ . In matrix form,

$$S^* := \begin{pmatrix} \frac{1}{2} & 1 & 1 & 1 & 1 & \cdots & 1 & 1 & \cdots \\ 0 & \frac{1}{2} & 1 & 1 & 1 & \cdots & 1 & 1 & \cdots \\ 0 & 0 & \frac{1}{2} & 1 & 1 & \cdots & 1 & 1 & \cdots \\ 0 & 0 & 0 & \frac{1}{2} & 1 & \cdots & 1 & 1 & \cdots \\ 0 & 0 & 0 & 0 & \frac{1}{2} & \cdots & 1 & 1 & \cdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \cdots & \cdots & \cdots \end{pmatrix}.$$

Moreover,  $\text{dom } S \subsetneq \text{dom } S^*$ ,  $S^* = -S$  on  $\text{dom } S$ , and  $S^*$  is not skew.

*Proof.* Let  $y = (y_n)_{n \in \mathbb{N}} \in \ell^2$  with  $\left( \sum_{i>n} y_i \right)_{n \in \mathbb{N}} \in \ell^2$ , and  $y^* = \left( \sum_{i>n} y_i + \frac{1}{2}y_n \right)_{n \in \mathbb{N}}$ . Now we show  $(y, y^*) \in \text{gra } S^*$ . Let  $s = \sum_{i \geq 1} y_i$  and  $x \in \text{dom } S$ . Then we have

$$\begin{aligned} \langle y, Sx \rangle + \langle y^*, -x \rangle &= \langle y, \frac{1}{2}x + \left( \sum_{i<n} x_i \right)_{n \in \mathbb{N}} \rangle + \langle \frac{1}{2}y + \left( \sum_{i>n} y_i \right)_{n \in \mathbb{N}}, -x \rangle \\ &= \langle y, \left( \sum_{i<n} x_i \right)_{n \in \mathbb{N}} \rangle + \langle \left( \sum_{i>n} y_i \right)_{n \in \mathbb{N}}, -x \rangle \\ &= \lim_n [y_2x_1 + y_3(x_1 + x_2) + \cdots + y_n(x_1 + \cdots + x_{n-1})] \\ &\quad - \lim_n [x_1(s - y_1) + x_2(s - y_1 - y_2) + \cdots + x_n(s - y_1 - \cdots - y_n)] \\ &= \lim_n [x_1(y_2 + \cdots + y_n) + x_2(y_3 + \cdots + y_n) + \cdots + x_{n-1}y_n] \\ &\quad - \lim_n [x_1(s - y_1) + x_2(s - y_1 - y_2) + \cdots + x_n(s - y_1 - \cdots - y_n)] \\ &= \lim_n [x_1(y_1 + y_2 + \cdots + y_n - s) + x_2(y_1 + y_2 + \cdots + y_n - s) + \cdots + x_n(y_1 + y_2 + \cdots + y_n - s)] \\ &= \lim_n [(x_1 + \cdots + x_n)(y_1 + y_2 + \cdots + y_n - s)] \\ &= 0. \end{aligned}$$

Hence  $(y, y^*) \in \text{gra } S^*$ .

On the other hand, let  $(a, a^*) \in \text{gra } S^*$  with  $a = (a_n)_{n \in \mathbb{N}}$  and  $a^* = (a_n^*)_{n \in \mathbb{N}}$ . Now we show

$$(13) \quad \left( \sum_{i>n} a_i \right)_{n \in \mathbb{N}} \in \ell^2 \text{ and } a^* = \left( \sum_{i>n} a_i + \frac{1}{2}a_n \right)_{n \in \mathbb{N}}.$$



Let  $e_n = (0, \dots, 0, 1, 0, \dots)$ : the  $n$ th entry is 1 and the others are 0. Then let  $y = -e_1 + e_n$ . Thus  $y \in \text{dom } S$  and  $Sy = (-\frac{1}{2}, -1, \dots, -1, -\frac{1}{2}, 0, \dots)$ . Then,

$$\begin{aligned} 0 &= \langle a^*, y \rangle + \langle -Sy, a \rangle = -a_1^* + a_n^* + \frac{1}{2}a_1 + \frac{1}{2}a_n + \sum_{i=2}^{n-1} a_i \\ (14) \quad &\Rightarrow a_n^* = a_1^* - \frac{1}{2}a_1 - \sum_{i=2}^{n-1} a_i - \frac{1}{2}a_n. \end{aligned}$$

Since  $a^* \in \ell^2$  and  $a \in \ell^2$ ,  $a_n^* \rightarrow 0, a_n \rightarrow 0$ . Thus by (14),

$$(15) \quad a_1^* = \frac{1}{2}a_1 + \sum_{i>1} a_i,$$

from which we see that  $\sum_{i \geq 1} a_i \in \mathbb{R}$ . Combining (14) and (15), we have

$$a_n^* = \sum_{i>n} a_i + \frac{1}{2}a_n$$

Thus, (13) holds. Hence (12) holds.

Now for  $x \in \text{dom } S$ , since  $\sum_{i \geq 1} x_i = 0$ , we have

$$\begin{aligned} S^*x &= \left( \frac{1}{2}x_n + \sum_{i>n} x_i \right)_{n \in \mathbb{N}} = \left( -\frac{1}{2}x_n + \sum_{i \geq n} x_i \right)_{n \in \mathbb{N}} \\ &= \left( -\frac{1}{2}x_n - \sum_{i<n} x_i \right)_{n \in \mathbb{N}} = -Sx. \end{aligned}$$

We note that  $S^*$  is not skew since for  $e_1 = (1, 0, \dots)$ ,  $\langle S^*e_1, e_1 \rangle = \langle 1/2e_1, e_1 \rangle = 1/2$ . As  $e_1 = (1, 0, 0, \dots, 0, \dots) \in \text{dom } S^*$  but  $e_1 \notin \text{dom } S$ . we have  $\text{dom } S \subsetneq \text{dom } S^*$ .  $\blacksquare$

**Proposition 3.7** *Let  $S$  be defined as in Example 3.1, let  $y = (y_1, y_2, \dots) \in \text{dom } S^*$ , and set  $s = \sum_{i \geq 1} y_i$ . Then*

$$(16) \quad \langle S^*y, y \rangle = \frac{1}{2}s^2.$$

*Proof.* By Proposition 3.6, we have  $s \in \mathbb{R}$  and

$$\begin{aligned}
\langle S^*y, y \rangle &= \left\langle \left( \sum_{i>n} y_i + \frac{1}{2}y_n \right)_{n \in \mathbb{N}}, y \right\rangle = \left\langle \left( \sum_{i \geq n} y_i - \frac{1}{2}y_n \right)_{n \in \mathbb{N}}, y \right\rangle \\
&= \lim_n [sy_1 + (s - y_1)y_2 + \cdots + (s - y_1 - y_2 - \cdots - y_{n-1})y_n - \frac{1}{2}(y_1^2 + y_2^2 + \cdots + y_n^2)] \\
&= \lim_n [s(y_1 + \cdots + y_n) - y_1y_2 - (y_1 + y_2)y_3 - \cdots - (y_1 + y_2 + \cdots + y_{n-1})y_n] \\
&\quad - \frac{1}{2} [y_1^2 + y_2^2 + \cdots + y_n^2] \\
&= \lim_n [s(y_1 + \cdots + y_n)] \\
&\quad - \lim_n [y_1y_2 + (y_1 + y_2)y_3 + \cdots + (y_1 + y_2 + \cdots + y_{n-1})y_n + \frac{1}{2}(y_1^2 + y_2^2 + \cdots + y_n^2)] \\
&= s^2 - \lim_n \frac{1}{2} [y_1 + y_2 + \cdots + y_n]^2 \\
&= s^2 - \frac{1}{2}s^2 \\
&= \frac{1}{2}s^2.
\end{aligned}$$

Hence (16) holds. ■

**Proposition 3.8** *Let  $S$  be defined as in Example 3.1. Then  $-S$  is not maximal monotone.*

*Proof.* By Proposition 3.2,  $-S$  is skew. Let  $e_1 = (1, 0, 0, \dots, 0, \dots)$ . Then  $e_1 \notin \text{dom } S = \text{dom}(-S)$ . Thus,  $(e_1, \frac{1}{2}e_1) \notin \text{gra}(-S)$ . We have for every  $y \in \text{dom } S$ ,

$$\langle e_1, \frac{1}{2}e_1 \rangle \geq 0 \text{ and } \langle e_1, -Sy \rangle + \langle y, \frac{1}{2}e_1 \rangle = -\frac{1}{2}y_1 + \frac{1}{2}y_1 = 0.$$

By Fact 3.4,  $(e_1, \frac{1}{2}e_1)$  is monotonically related to  $\text{gra}(-S)$ . We deduce that  $-S$  is not maximal monotone. ■

We proceed to show that for every maximal monotone and skew operator  $S$ , the operator  $-S$  has a unique maximal monotone extension, namely  $S^*$ .

**Theorem 3.9** *Let  $S : \text{dom } S \rightarrow X$  be a maximal monotone skew operator. Then  $-S$  has a unique maximal monotone extension:  $S^*$ .*

*Proof.* By Fact 2.2,  $\text{gra}(-S) \subseteq \text{gra } S^*$ . Assume  $T$  is a maximal monotone extension of  $-S$ . Let  $(x, x^*) \in \text{gra } T$ . Then  $(x, x^*)$  is monotonically related to  $\text{gra}(-S)$ . By Fact 3.4,

$$\langle x^*, y \rangle + \langle -x, Sy \rangle = \langle x^*, y \rangle + \langle x, -Sy \rangle = 0, \quad \forall y \in \text{dom } S.$$

Thus  $(x, x^*) \in \text{gra } S^*$ . Since  $(x, x^*) \in \text{gra } T$  is arbitrary, we have  $\text{gra } T \subseteq \text{gra } S^*$ . By Fact 2.3,  $S^*$  is maximal monotone. Hence  $T = S^*$ . ■

**Remark 3.10** Note that [23, Proposition 17] also implies that  $-S$  has a unique maximal monotone extension, where  $S$  is as in Theorem 3.9.

**Remark 3.11** Define the *right and left shift operators*  $R, L : \ell^2 \rightarrow \ell^2$  by

$$Rx = (0, x_1, x_2, \dots), \quad Lx = (x_2, x_3, \dots), \quad \forall x = (x_1, x_2, \dots) \in \ell^2.$$

One can verify that in Example 3.1

$$S = (\text{Id} - R)^{-1} - \frac{\text{Id}}{2}, \quad S^* = (\text{Id} - L)^{-1} - \frac{\text{Id}}{2}.$$

The maximal monotone operators  $(\text{Id} - R)^{-1}$  and  $(\text{Id} - L)^{-1}$  have been utilized by Phelps and Simons, see [14, Example 7.4].

### 3.2 An answer to Svaiter's question

**Definition 3.12** Let  $S : X \rightrightarrows X$  be skew. We say  $S$  is maximal skew (termed “maximal self-cancelling” in [21]) if no proper enlargement (in the sense of graph inclusion) of  $S$  is skew. We say  $T$  is a maximal skew extension of  $S$  if  $T$  is maximal skew and  $\text{gra} T \supseteq \text{gra} S$ .

**Lemma 3.13** Let  $S : X \rightrightarrows X$  be a maximal monotone skew operator. Then both  $S$  and  $-S$  are maximal skew.

*Proof.* Clearly,  $S$  is maximal skew. Now we show  $-S$  is maximal skew. Let  $T$  be a skew operator such that  $\text{gra}(-S) \subseteq \text{gra} T$ . Thus,  $\text{gra} S \subseteq \text{gra}(-T)$ . Since  $-T$  is monotone and  $S$  is maximal monotone,  $\text{gra} S = \text{gra}(-T)$ . Then  $-S = T$ . Hence  $-S$  is maximal skew. ■

**Fact 3.14 (Svaiter)** (See [21].) Let  $S : X \rightrightarrows X$  be maximal skew. Then either  $-S^*$  (i.e.,  $S^\perp$ ) or  $S^*$  (i.e.,  $-S^\perp$ ) is maximal monotone.

In [21], Svaiter asked whether or not  $-S^*$  (i.e.,  $S^\perp$ ) is maximal monotone if  $S$  is maximal skew. Now we can give a negative answer, even though  $S$  is maximal monotone and skew.

**Theorem 3.15** Let  $S$  be defined as in Example 3.1. Then  $S$  is maximal skew, but  $-S^*$  is not monotone, so not maximal monotone.

*Proof.* Let  $e_1 = (1, 0, 0, \dots, 0, \dots)$ . By Proposition 3.6,  $(e_1, -\frac{1}{2}e_1) \in \text{gra}(-S^*)$ , but  $\langle e_1, -\frac{1}{2}e_1 \rangle = -\frac{1}{2} < 0$ . Hence  $-S^*$  is not monotone. ■

By Theorem 3.15,  $-S^*$  (i.e.,  $S^\perp$ ) is not always maximal monotone. Can one improve Svaiter's result: “If  $S$  is maximal skew, then  $S^*$  (i.e.,  $-S^\perp$ ) is always maximal monotone?”

**Theorem 3.16** There exists a maximal skew operator  $T$  on  $\ell^2$  such that  $T^*$  is not maximal monotone. Consequently, Svaiter's result is optimal.

*Proof.* Let  $T = -S$ , where  $S$  is defined as in Example 3.1. By Lemma 3.13,  $T$  is maximal skew. Then by Theorem 3.15 and Fact 2.1(iii),  $T^* = (-S)^* = -S^*$  is not maximal monotone. Hence Svaiter's result cannot be further improved. ■

### 3.3 The maximal monotonicity and Fitzpatrick functions of a sum

**Example 3.17** ( *$S + S^*$  fails to be maximal monotone*) Let  $S$  be defined in Example 3.1. Then neither  $S$  nor  $S^*$  has full domain. By Fact 2.2,  $\forall x \in \text{dom}(S + S^*) = \text{dom } S$ , we have

$$(S + S^*)x = 0.$$

Thus  $S + S^*$  has a proper monotone extension from  $\text{dom}(S + S^*)$  to the 0 map on  $X$ . Consequently,  $S + S^*$  is not maximal monotone. This supplies a different example for showing that the constraint qualification in the sum problem of maximal monotone operators can not be substantially weakened, see [14, Example 7.4].

We now compute  $F_S, F_{S^*}, F_{S+S^*}$ . As a result, we see that  $F_{S+S^*} \neq F_S \square_2 F_{S^*}$  even though  $S, S^*$  are maximal monotone with  $\text{dom } S - \text{dom } S^*$  being dense in  $\ell^2$ . Since  $\text{ran}(S_+ + (S^*)_+) = \{0\}$  and  $F_{S+S^*} \neq F_S \square_2 F_{S^*}$ , this also means that Fact 2.8(i) fails for discontinuous linear maximal monotone operators.

**Lemma 3.18** *Let  $S : \text{dom } S \rightarrow X$  be a maximal monotone skew linear operator. Then*

$$\begin{aligned} F_S &= \iota_{\text{gra}(-S^*)}, \\ F_{S^*}^{\text{T}} &= F_{S^*} = \iota_{\text{gra } S^*} + \langle \cdot, \cdot \rangle. \end{aligned}$$

*Proof.* By [5, Proposition 5.5],

$$F_S^* = (\iota_{\text{gra } S})^{\text{T}}.$$

Then

$$(17) \quad F_S = (F_S^{\text{T}})^{* \text{T}} = (\iota_{\text{gra } S})^{* \text{T}} = (\iota_{\text{gra } S}^{\text{T}})^* = (\iota_{\text{gra } S^{-1}})^* = \iota_{(\text{gra } S^{-1})^\perp} = \iota_{\text{gra}(-S^*)}.$$

From Fact 2.2,  $\text{gra } -S \subseteq \text{gra } S^*$ , we have

$$F_{S^*} \geq F_{-S} = \iota_{\text{gra } -(-S)^*} = \iota_{\text{gra } S^*},$$

this shows that  $\text{dom } F_{S^*} \subseteq \text{gra } S^*$ . By Fact 2.4,  $F_{S^*}(x, x^*) = \langle x, x^* \rangle \forall (x, x^*) \in \text{gra } S^*$ . Hence  $F_{S^*} = \iota_{\text{gra } S^*} + \langle \cdot, \cdot \rangle$ . Again by [5, Proposition 5.5],  $F_{S^*}^{\text{T}} = \iota_{\text{gra } S^*} + \langle \cdot, \cdot \rangle$ .  $\blacksquare$

**Theorem 3.19** *Let  $S$  be defined as in Example 3.1. Then*

$$(18) \quad \begin{aligned} F_{S+S^*}(x, x^*) &= \iota_{X \times \{0\}}(x, x^*) \\ F_S \square_2 F_{S^*}(x, x^*) &= \begin{cases} \frac{1}{2}s^2, & \text{if } (x, x^*) \in \text{dom } S^* \times \{0\} \text{ with } s = \sum_{i \geq 1} x_i; \\ \infty & \text{otherwise.} \end{cases} \end{aligned}$$

Consequently,  $F_S \square_2 F_{S^*} \neq F_{S+S^*}$ .

*Proof.* By Fact 2.2,

$$(19) \quad (S + S^*)|_{\text{dom } S} = 0.$$

Let  $(x, x^*) \in X \times X$ . Using (19) and Fact 2.2, we have

$$(20) \quad F_{S+S^*}(x, x^*) = \sup_{a \in \text{dom } S} \langle x^*, a \rangle = \iota_{(\text{dom } S)^\perp}(x^*) = \iota_{\{0\}}(x^*) = \iota_{X \times \{0\}}(x, x^*).$$

Then by Fact 2.7, we have

$$(21) \quad F_S \square_2 F_{S^*}(x, x^*) = \infty, \quad x^* \neq 0.$$

It follows from Lemma 3.18 that

$$(22) \quad \begin{aligned} F_S \square_2 F_{S^*}(x, 0) &= \inf_{y^* \in X} \{F_S(x, y^*) + F_{S^*}(x, -y^*)\} \\ &= \inf_{y^* \in X} \{\iota_{\text{gra}(-S^*)}(x, y^*) + \iota_{\text{gra } S^*}(x, -y^*) + \langle x, -y^* \rangle\} \\ &= \inf_{y^* \in X} \{\iota_{\text{gra } S^*}(x, -y^*) + \langle x, -y^* \rangle\}. \end{aligned}$$

Thus,  $F_S \square_2 F_{S^*}(x, 0) = \infty$  if  $x \notin \text{dom } S^*$ . Now suppose  $x \in \text{dom } S^*$  and  $s = \sum_{i \geq 1} x_i$ . Then by (22) and Proposition 3.7, we have

$$F_S \square_2 F_{S^*}(x, 0) = \langle x, S^* x \rangle = \frac{1}{2} s^2.$$

Combine the results above, (18) holds. Since  $\text{dom } S^* \neq X$ ,  $F_S \square_2 F_{S^*} \neq F_{S+S^*}$ . ■

**Remark 3.20** [5, Theorem 7.6] shows that: Let  $A : X \rightrightarrows X$  be a maximal monotone linear relation. Then  $A^* = -A$  if and only if  $\text{dom } A = \text{dom } A^*$  and  $F_A = F_A^{*\top}$ . Let  $A = S^*$  with  $S$  defined as in Example 3.1. Lemma 3.18 shows that  $F_A = F_A^{*\top}$ , but  $A^* = S \neq -S^* = -A$ . Hence the requirement  $\text{dom } A = \text{dom } A^*$  can not be omitted.

## 4 The inverse Volterra operator on $L^2[0, 1]$

Let  $V$  be the Volterra integral operator. In this section, we systematically study  $T = V^{-1}$  and its skew part  $S := \frac{1}{2}(T - T^*)$ . It turns out that  $T$  is neither skew nor symmetric and that its skew part  $S$  admits two maximal monotone and skew extensions  $T_1, T_2$  (in fact, anti-self-adjoint) even though  $\text{dom } S$  is a dense linear subspace of  $L^2[0, 1]$ . This will give another simpler example of Phelps-Simons' showing that the constraint qualification for the sum of monotone operators cannot be significantly weakened, see [19, Theorem 5.5] or [22]. We compute the Fitzpatrick functions  $F_T, F_{T^*}, F_{T+T^*}$ , and we show that  $F_T \square_2 F_{T^*} \neq F_{T+T^*}$ . This shows that the constraint qualification for the formula of the Fitzpatrick function of the sum of two maximal monotone operators cannot be significantly weakened either.

**Definition 4.1** ([5]) *Let  $T : X \rightrightarrows X$  be a linear relation. We say that  $T$  is symmetric if  $\text{gra } T \subseteq \text{gra } T^*$ ;  $T$  is self-adjoint if  $T^* = T$  and anti-self-adjoint if  $T^* = -T$ .*

## 4.1 Properties of the Volterra operator and its inverse

To study the Volterra operator and its inverse, we shall frequently need the following generalized integration-by-parts formula, see [20, Theorem 6.90].

**Fact 4.2 (Generalized integration by parts)** *Assume that  $x, y$  are absolutely continuous functions on the interval  $[a, b]$ . Then*

$$\int_a^b xy' + \int_a^b x'y = x(b)y(b) - x(a)y(a).$$

Fact 2.3 allows us to claim that

**Proposition 4.3** *Let  $A : X \rightrightarrows X$  be a linear relation. If  $A^* = -A$ , then both  $A$  and  $-A$  are maximal monotone and skew.*

*Proof.* Since  $A = -A^*$ , we have that  $\text{dom } A = \text{dom } A^*$  and that  $A$  has closed graph. Now  $\forall x \in \text{dom } A$ , by Fact 2.1(iv),

$$\langle Ax, x \rangle = \langle x, A^*x \rangle = -\langle x, Ax \rangle \quad \Rightarrow \quad \langle Ax, x \rangle = 0.$$

Hence  $A$  and  $-A$  are skew. As  $A^* = -A$  is monotone, Fact 2.3 shows that  $A$  is maximal monotone.

Now  $-A = A^* = -(-A)^*$  and  $-A$  is a linear relation. Similar arguments show that  $-A$  is maximal monotone. ■

**Example 4.4 (Volterra operator)** (See [2, Example 3.3].) Set  $X = L^2[0, 1]$ . The *Volterra integration operator* [13, Problem 148] is defined by

$$(23) \quad V : X \rightarrow X : x \mapsto Vx, \quad \text{where} \quad Vx : [0, 1] \rightarrow \mathbb{R} : t \mapsto \int_0^t x,$$

and its adjoint is given by

$$t \mapsto (V^*x)(t) = \int_t^1 x, \quad \forall x \in X.$$

Then

(i) Both  $V$  and  $V^*$  are maximal monotone since they are monotone, continuous and linear.

(ii) Both ranges

$$(24) \quad \text{ran } V = \{x \in L^2[0, 1] : x \text{ is absolutely continuous, } x(0) = 0, x' \in L^2[0, 1]\},$$

and

$$(25) \quad \text{ran } V^* = \{x \in L^2[0, 1] : x \text{ is absolutely continuous, } x(1) = 0, x' \in L^2[0, 1]\},$$

are dense in  $X$ , and both  $V$  and  $V^*$  are one-to-one.

(iii)  $\text{ran } V \cap \text{ran } V^* = \{Vx \mid x \in e^\perp\}$ , where  $e \equiv 1 \in L^2[0, 1]$ .

(iv) Define  $V_+x := \frac{1}{2}(V + V^*)(x) = \frac{1}{2}\langle e, x \rangle e$ . Then  $V_+$  is self-adjoint and

$$\text{ran } V_+ = \text{span}\{e\}.$$

(v) Define  $V_\circ x := \frac{1}{2}(V - V^*)(x) : t \mapsto \frac{1}{2}[\int_0^t x - \int_t^1 x] \quad \forall x \in L^2[0, 1], t \in [0, 1]$ . Then  $V_\circ$  is anti-self-adjoint and

$$\text{ran } V_\circ = \{x \in L^2[0, 1] : x \text{ is absolutely continuous on } [0, 1], x' \in L^2[0, 1], x(0) = -x(1)\}.$$

*Proof.* (i) By Fact 4.2,

$$\langle x, Vx \rangle = \int_0^1 x(t) \int_0^t x(s) ds dt = \frac{1}{2} \left( \int_0^1 x(s) ds \right)^2 \geq 0,$$

so  $V$  is monotone.

As  $\text{dom } V = L^2[0, 1]$  and  $V$  is continuous,  $\text{dom } V^* = L^2[0, 1]$ . Let  $x, y \in L^2[0, 1]$ . We have

$$\begin{aligned} \langle Vx, y \rangle &= \int_0^1 \int_0^t x(s) ds y(t) dt = \int_0^1 x(t) dt \int_0^1 y(s) ds - \int_0^1 \int_0^t y(s) ds x(t) dt \\ &= \int_0^1 \left( \int_0^1 y(s) ds - \int_0^t y(s) ds \right) x(t) dt = \int_0^1 \int_t^1 y(s) ds x(t) dt = \langle V^*y, x \rangle, \end{aligned}$$

thus  $(V^*y)(t) = \int_t^1 y(s) ds \quad \forall t \in [0, 1]$ .

(ii) To show (24), if  $z \in \text{ran } V$ , then

$$z(t) = \int_0^t x \quad \text{for some } x \in L^2[0, 1],$$

and hence  $z(0) = 0$ ,  $z$  is absolutely continuous, and  $z' = x \in L^2[0, 1]$ . On the other hand, if  $z(0) = 0$ ,  $z$  is absolutely continuous,  $z' \in L^2[0, 1]$ , then  $z = Vz'$ .

To show (25), if  $z \in \text{ran } V^*$ , then

$$z(t) = \int_t^1 x \quad \text{for some } x \in L^2[0, 1],$$

and hence  $z(1) = 0$ ,  $z$  is absolutely continuous, and  $z' = -x \in L^2[0, 1]$ . On the other hand, if  $z(1) = 0$ ,  $z$  is absolutely continuous,  $z' \in L^2[0, 1]$ , then  $z = V^*(-z')$ .

(iii) follows from (ii) (or see [2]).

(iv) is clear.

(v) If  $x$  is absolutely continuous,  $x(0) = -x(1)$ ,  $x' \in L^2[0, 1]$ , we have

$$V_{\circ}x'(t) = \frac{1}{2} \left( \int_0^t x' - \int_t^1 x' \right) = \frac{1}{2} \left( x(t) - x(0) - x(1) + x(t) \right) = x(t).$$

This shows that  $x \in \text{ran } V_{\circ}$ . Conversely, if  $x \in \text{ran } V_{\circ}$ , i.e.,

$$x(t) = \frac{1}{2} \int_0^t y - \frac{1}{2} \int_t^1 y \quad \text{for some } y \in L^2[0, 1],$$

then  $x$  is absolutely continuous,  $x' = y \in L^2[0, 1]$  and  $x(0) = -x(1) = -\frac{1}{2} \int_0^1 y$ . ■

**Theorem 4.5 (Inverse Volterra operator=Differentiation operator)** *Let  $X = L^2[0, 1]$ , and  $V$  be the Volterra integration operator. We let  $T = V^{-1}$  and  $D = \text{dom } T \cap \text{dom } T^*$ . Then the following hold.*

(i)  $T : \text{dom } T \rightarrow X$  is given by  $Tx = x'$  with

$$\text{dom } T = \{x \in L^2[0, 1] : x \text{ is absolutely continuous, } x(0) = 0, x' \in L^2[0, 1]\},$$

and  $T^* : \text{dom } T^* \rightarrow X$  is given by  $T^*x = -x'$  with

$$\text{dom } T^* = \{x \in L^2[0, 1] : x \text{ is absolutely continuous, } x(1) = 0, x' \in L^2[0, 1]\}.$$

Both  $T$  and  $T^*$  are maximal monotone linear operators.

(ii)  $T$  is neither skew nor symmetric.

(iii) The linear subspace

$$D = \{x \in L^2[0, 1] : x \text{ is absolutely continuous, } x(0) = x(1) = 0, x' \in L^2[0, 1]\}$$

is dense in  $X$ . Moreover,  $T$  and  $T^*$  are skew on  $D$ .

*Proof.* (i):  $T$  and  $T^*$  are maximal monotone because  $T = V^{-1}$ , and  $T^* = (V^{-1})^* = (V^*)^{-1}$  and Example 4.4(i). By Example 4.4(ii),  $T : L^2[0, 1] \rightarrow L^2[0, 1]$  has

$$\begin{aligned} \text{dom } T &= \{x \in L^2[0, 1] : x \text{ is absolutely continuous, } x(0) = 0, x' \in L^2[0, 1]\} \\ \text{dom } T^* &= \{x \in L^2[0, 1] : x \text{ is absolutely continuous, } x(1) = 0, x' \in L^2[0, 1]\} \\ Tx &= x', \quad \forall x \in \text{dom } T, \quad T^*y = -y' \text{ and } \forall y \in \text{dom } T^*. \end{aligned}$$

Note that by Fact 4.2,

$$(26) \quad \langle Tx, x \rangle = \int_0^1 x'x = \frac{1}{2}x^2(1) - \frac{1}{2}x^2(0) = \frac{1}{2}x(1)^2 \quad \forall x \in \text{dom } T,$$

$$(27) \quad \langle T^*x, x \rangle = \int_0^1 -x'x = -\left(\frac{1}{2}x(1)^2 - \frac{1}{2}x(0)^2\right) = \frac{1}{2}x(0)^2 \quad \forall x \in \text{dom } T^*.$$



(ii): Letting  $x(t) = t, y(t) = t^2$  we have

$$\langle Tx, x \rangle = \int_0^1 t = \frac{1}{2}, \quad \langle x, Ty \rangle = \int_0^1 2t^2 = \frac{2}{3} \neq \frac{1}{3} = \int_0^1 t^2 = \langle Tx, y \rangle \quad \Rightarrow \langle Tx, x \rangle \neq 0, \langle Tx, y \rangle \neq \langle x, Ty \rangle.$$

(iii): By (i),  $D = \text{dom } T \cap \text{dom } T^*$  is clearly a linear subspace. For  $x \in D, x(0) = x(1) = 0$ , from (26) and (27),

$$\langle Tx, x \rangle = \frac{1}{2}x(1)^2 = 0, \quad \langle T^*x, x \rangle = \frac{1}{2}x(0)^2 = 0.$$

Hence both  $T$  and  $T^*$  are skew on  $D$ . The fact that  $D$  is dense in  $L^2[0, 1]$  follows from [20, Theorem 6.111].  $\blacksquare$

Our proof of (ii), (iii) in the following theorem follows the ideas of [17, Example 13.4].

**Theorem 4.6 (The skew part of inverse Volterra operator)** *Let  $X = L^2[0, 1]$ , and  $T$  be defined as in Theorem 4.5. Let  $S := \frac{T-T^*}{2}$ .*

(i)  $Sx = x' (\forall x \in \text{dom } S)$  and  $\text{gra } S = \{(Vx, x) \mid x \in e^\perp\}$ , where  $e \equiv 1 \in L^2[0, 1]$ . In particular,

$$\begin{aligned} \text{dom } S &= \{x \in L^2[0, 1] : x \text{ is absolutely continuous, } x(0) = x(1) = 0, x' \in L^2[0, 1]\}, \\ \text{ran } S &= \{y \in L^2[0, 1] : \langle e, y \rangle = 0\} = e^\perp. \end{aligned}$$

Moreover,  $\text{dom } S$  is dense, and

$$(28) \quad S^{-1} = V|_{e^\perp}, \quad (-S)^{-1} = V^*|_{e^\perp},$$

consequently,  $S$  is skew, and neither  $S$  nor  $-S$  is maximal monotone.

(ii) The adjoint of  $S$  has  $\text{gra } S^* = \{(V^*x^* + le, x^*) \mid x^* \in X, l \in \mathbb{R}\}$ . More precisely,

$$\begin{aligned} S^*x &= -x' \quad \forall x \in \text{dom } S^*, \text{ with} \\ \text{dom } S^* &= \{x \in L^2[0, 1] : x \text{ is absolutely continuous on } [0, 1], x' \in L^2[0, 1]\}, \\ \text{ran } S^* &= L^2[0, 1]. \end{aligned}$$

Neither  $S^*$  nor  $-S^*$  is monotone. Moreover,  $S^{**} = S$ .

(iii) Let  $T_1 : \text{dom } T_1 \rightarrow X$  be defined by

$$T_1x = x', \quad \forall x \in \text{dom } T_1 := \{x \in L^2[0, 1] : x \text{ is absolutely continuous, } x(0) = x(1), x' \in L^2[0, 1]\}.$$

Then  $T_1^* = -T_1$ ,

$$(29) \quad \text{ran } T_1 = e^\perp.$$

Hence  $T_1$  is skew, and a maximal monotone extension of  $S$ ; and  $-T_1$  is skew and a maximal monotone extension of  $-S$ .

*Proof.* (i): By Theorem 4.5(iii), we directly get  $\text{dom } S$ . Now  $(\forall x \in \text{dom } S = \text{dom } T \cap \text{dom } T^*) Tx = x'$  and  $T^*x = -x'$ , so  $Sx = x'$ . Then Example 4.4(iii) implies  $\text{gra } S = \{(Vx, x) \mid x \in e^\perp\}$ . Hence

$$(30) \quad \text{gra } S^{-1} = \{(x, Vx) : x \in e^\perp\}.$$

Theorem 4.5(iii) implies  $\text{dom } S$  is dense. Furthermore,  $\text{gra}(-S) = \{(Vx, -x) : x \in e^\perp\}$ , so

$$\text{gra}(-S)^{-1} = \{(x, -Vx) : x \in e^\perp\}.$$

Since

$$V^*x(t) = \int_t^1 x - 0 = \int_t^1 x - \int_0^1 x = - \int_0^t x = -Vx(t) \quad \forall t \in [0, 1], \forall x \in e^\perp$$

we have  $-Vx = V^*x, \forall x \in e^\perp$ . Then

$$(31) \quad \text{gra}(-S)^{-1} = \{(x, V^*x) : x \in e^\perp\}.$$

Hence, (30) and (31) together establish (28). As both  $V, V^*$  are maximal monotone with full domain, we conclude that  $S^{-1}, (-S)^{-1}$  are not maximal monotone, thus  $S, -S$  are not maximal monotone.

(ii): By (i), we have

$$\begin{aligned} (x, x^*) \in \text{gra } S^* &\Leftrightarrow \langle -x, y \rangle + \langle x^*, Vy \rangle = 0, \quad \forall y \in e^\perp \\ &\Leftrightarrow \langle -x + V^*x^*, y \rangle = 0, \quad \forall y \in e^\perp \Leftrightarrow x - V^*x^* \in \text{span}\{e\}. \end{aligned}$$

Equivalently,  $x = V^*x^* + ke$  for some  $k \in \mathbb{R}$ . This means that  $x$  is absolutely continuous,  $x^* = -x' \in L^2[0, 1]$ . On the other hand, if  $x$  is absolutely continuous and  $x' \in L^2[0, 1]$ , observe that

$$x(t) = \int_t^1 -x' + x(1)e,$$

so that  $x - V^*(-x') \in \text{span}\{e\}$  and  $(x, -x') \in \text{gra } S^*$ . It follows that

$$\text{dom } S^* = \{x \in L^2[0, 1] : x \text{ is absolutely continuous on } [0, 1], x' \in L^2[0, 1]\},$$

$$\text{ran } S^* = L^2[0, 1], \quad \text{and}$$

$$S^*x = -x', \quad \forall x \in \text{dom } S^*.$$

Since

$$\langle S^*x, x \rangle = - \int_0^1 x'x = - \left( \frac{1}{2}x(1)^2 - \frac{1}{2}x(0)^2 \right),$$

we conclude that neither  $S^*$  nor  $-S^*$  is monotone.

We proceed to show that  $S^{**} = S$ . Note that  $\forall x \in \text{dom } S^*, z \in \text{dom } S$ , we have  $z(0) = z(1) = 0$  and

$$\langle S^*x, z \rangle = \int_0^1 -x'z = - \left( x(1)z(1) - x(0)z(0) - \int_0^1 xz' \right) = \int_0^1 xz' = \langle x, Sz \rangle,$$

this implies that  $S^{**}z = Sz$ ,  $\forall z \in \text{dom } S$ , i.e.,  $S^{**}|_{\text{dom } S} = S$ . Suppose now that  $x \in \text{dom } S^{**}$ ,  $\varphi = S^{**}x$ . Put  $\Phi = V$ . Then  $\forall z \in \text{dom } S^*$ ,

$$\begin{aligned}\langle S^*z, x \rangle &= \int_0^1 -z'x = \langle z, S^{**}x \rangle \\ &= \langle z, \varphi \rangle = \int_0^1 z\varphi = [z(1)\Phi(1) - z(0)\Phi(0)] - \int_0^1 \Phi z' \\ &= z(1)\Phi(1) - \int_0^1 \Phi z'.\end{aligned}$$

Using  $z = e \in \text{dom } S^*$  gives  $\Phi(1) = 0$ . It follows that

$$\int_0^1 [\Phi - x]z' = 0, \quad \forall z \in \text{dom } S^* \Rightarrow \Phi - x \in (\text{ran } S^*)^\perp,$$

then  $\Phi = x$  since  $\text{ran } S^* = L^2[0, 1]$ . As  $\Phi(1) = \Phi(0) = 0$  and  $\Phi$  is absolutely continuous, we have  $x \in \text{dom } S$ . Since  $x \in \text{dom } S^{**}$  was arbitrary, we conclude that  $\text{dom } S^{**} \subseteq \text{dom } S$ . Hence  $S^{**} = S$ . (Alternatively,  $V$  is continuous  $\Rightarrow V|_{e^\perp}$  has closed graph  $\Rightarrow S^{-1}$  has closed graph  $\Rightarrow S$  has closed graph  $\Rightarrow \text{gra } S = \text{gra } S^{**} \Rightarrow S^{**} = S$ .)

(iii): To show (29), suppose that  $x$  is absolutely continuous and that  $x(0) = x(1)$ . Then

$$\int_0^1 x' = x(1) - x(0) = 0 \quad \Rightarrow T_1x = x' \in e^\perp.$$

Conversely, if  $x \in L^2[0, 1]$  satisfies  $\langle e, x \rangle = 0$ , we define  $z = Vx$ , then  $z$  is absolutely continuous,  $z(0) = z(1)$ ,  $T_1z = x$ . Hence  $\text{ran } T_1 = e^\perp$ .

$T_1$  is skew, because for every  $x \in \text{dom } T_1$ , we have

$$\langle T_1x, x \rangle = \int_0^1 x'x = \frac{1}{2}x(1)^2 - \frac{1}{2}x(0)^2 = 0.$$

Moreover,  $T_1^* = -T_1$ : indeed, as  $T_1$  is skew, by Fact 2.2,  $\text{gra}(-T_1) \subseteq \text{gra } T_1^*$ . To show that  $T_1^* = -T_1$ , take  $z \in \text{dom } T_1^*$ ,  $\varphi = T_1^*z$ . Put  $\Phi = V\varphi$ . We have  $\forall y \in \text{dom } T_1$ ,

$$(32) \quad \int_0^1 y'z = \langle T_1y, z \rangle = \langle T_1^*z, y \rangle = \langle \varphi, y \rangle = \int_0^1 y\varphi = \int_0^1 y\Phi'$$

$$(33) \quad = [\Phi(1)y(1) - \Phi(0)y(0)] - \int_0^1 \Phi y'.$$

Using  $y = e \in \text{dom } T_1$  gives  $\Phi(1) - \Phi(0) = 0$ , from which  $\Phi(1) = \Phi(0) = 0$ . It follows from (32)–(33) that  $\int_0^1 y'(z + \Phi) = 0 \forall y \in \text{dom } T_1$ . Since  $\text{ran } T_1 = e^\perp$ ,  $z + \Phi \in \text{span}\{e\}$ , say  $z + \Phi = ke$  for some constant  $k \in \mathbb{R}$ . Then  $z$  is absolutely continuous,  $z(0) = z(1)$  since  $\Phi(0) = \Phi(1) = 0$ , and  $T_1^*z = \varphi = \Phi' = -z'$ . This implies that  $\text{dom } T_1^* \subseteq \text{dom } T_1$ . Then by Fact 2.2,  $T_1^* = -T_1$ . It remains to apply Proposition 4.3. ■

**Fact 4.7** Let  $A : X \rightrightarrows X$  be a multifunction. Then  $(-A)^{-1} = A^{-1} \circ (-\text{Id})$ . If  $A$  is a linear relation, then

$$(-A)^{-1} = -A^{-1}.$$

*Proof.* This follows from the set-valued inverse definition. Indeed,  $x \in (-A)^{-1}(x^*) \Leftrightarrow (x, x^*) \in \text{gra}(-A) \Leftrightarrow (x, -x^*) \in \text{gra} A \Leftrightarrow x \in A^{-1}(-x^*)$ . When  $A$  is a linear relation,  $x \in (-A)^{-1}(x^*) \Leftrightarrow (x, -x^*) \in \text{gra} A \Leftrightarrow (-x, x^*) \in \text{gra} A \Leftrightarrow -x \in A^{-1}x^* \Leftrightarrow x \in -A^{-1}(x^*)$ . ■

**Theorem 4.8 (The inverse of the skew part of Volterra operator)** Let  $X = L^2[0, 1]$ , and  $V$  be the Volterra integration operator, and  $V_\circ : L^2[0, 1] \rightarrow L^2[0, 1]$  be given by

$$V_\circ = \frac{V - V^*}{2}.$$

Define  $T_2 : \text{dom} T_2 \rightarrow L^2[0, 1]$  by  $T_2 = V_\circ^{-1}$ . Then

$$(i) \quad T_2 x = x', \quad \forall x \in \text{dom} T_2 \text{ where} \\ (34)$$

$$\text{dom} T_2 = \{x \in L^2[0, 1] : x \text{ is absolutely continuous on } [0, 1], x' \in L^2[0, 1], x(0) = -x(1)\}.$$

$$(ii) \quad T_2^* = -T_2, \text{ and both } T_2, -T_2 \text{ are maximal monotone and skew.}$$

*Proof.* (i) Since

$$V_\circ x(t) = \frac{1}{2} \left( \int_0^t x - \int_t^1 x \right),$$

$V_\circ$  is a one-to-one map. Then

$$V_\circ^{-1} \left( \frac{1}{2} \left( \int_0^t x - \int_t^1 x \right) \right) = x(t) = \left( \frac{1}{2} \left( \int_0^t x - \int_t^1 x \right) \right)',$$

which implies  $T_2 x = V_\circ^{-1} x = x'$  for  $x \in \text{ran} V_\circ$ . As  $\text{dom} T_2 = \text{ran} V_\circ$ , by Example 4.4(v),  $\text{ran} V_\circ$  can be written as (34).

(ii) Since  $\text{dom} V = \text{dom} V^* = L^2[0, 1]$ ,  $V_\circ$  is skew on  $L^2[0, 1]$ , so maximal monotone. Then  $T_2 = V_\circ^{-1}$  is maximal monotone.

Since  $V_\circ$  is skew and  $\text{dom} V_\circ = L^2[0, 1]$ , we have  $V_\circ^* = -V_\circ$ , by Fact 4.7,

$$T_2^* = (V_\circ^{-1})^* = (V_\circ^*)^{-1} = (-V_\circ)^{-1} = -V_\circ^{-1} = -T_2.$$

By Proposition 4.3, both  $T_2$  and  $-T_2$  are maximal monotone and skew. ■

**Remark 4.9** Note that while  $V_\circ$  is continuous on  $L^2[0, 1]$ , the operator  $S$  given in Example 3.1 is discontinuous.

Combining Theorem 4.5, Theorem 4.6 and Theorem 4.8, we can summarize the nice relationships among the differentiation operators encountered in this section.

**Corollary 4.10** *The domain of the skew operator  $S$  is dense in  $L^2[0, 1]$ . Neither  $S$  nor  $-S$  is maximal monotone. Neither  $S^*$  nor  $-S^*$  is monotone.*

*The linear operators  $S, T, T_1, T_2$  satisfy:*

$$\text{gra } S \subsetneq \text{gra } T \subsetneq \text{gra}(-S^*),$$

$$\text{gra } S \subsetneq \text{gra } T_1 \subsetneq \text{gra}(-S^*),$$

$$\text{gra } S \subsetneq \text{gra } T_2 \subsetneq \text{gra}(-S^*).$$

*While  $S$  is skew,  $T, T_1, T_2$  are maximal monotone and  $T_1, T_2$  are skew. Also,*

$$\text{gra}(-S) \subsetneq \text{gra}(T^*) \subsetneq \text{gra } S^*,$$

$$\text{gra}(-S) \subsetneq \text{gra}(-T_1) \subsetneq \text{gra } S^*,$$

$$\text{gra}(-S) \subsetneq \text{gra}(-T_2) \subsetneq \text{gra } S^*.$$

*While  $-S$  is skew,  $T^*, -T_1, -T_2$  are maximal monotone and  $-T_1, -T_2$  are skew.*

**Remark 4.11** (i). Note that while  $T_1, T_2$  are maximal monotone,  $-T_1, -T_2$  are also maximal monotone. This is in stark contrast with the maximal monotone skew operator given in Proposition 3.5 and Proposition 3.8 such that its negative is not maximal monotone.

(ii). Even though the skew operator  $S$  in Theorem 4.6 has  $\text{dom } S$  dense in  $L^2[0, 1]$ , it still admits two distinct maximal monotone and skew extensions  $T_1, T_2$ .

## 4.2 Consequences on sum of maximal monotone operators and Fitzpatrick functions of a sum

**Example 4.12** ( $T + T^*$  fails to be maximal monotone) Let  $T$  be defined as in Theorem 4.5. Now  $\forall x \in \text{dom } T \cap \text{dom } T^*$ , we have

$$Tx + T^*x = x' - x' = 0.$$

Thus  $T + T^*$  has a proper monotone extension from  $\text{dom } T \cap \text{dom } T^* \subsetneq X$  to the 0 map on  $X$ . Consequently,  $T + T^*$  is not maximal monotone. Note that  $\text{dom } T \cap \text{dom } T^*$  is dense in  $X$  and that  $\text{dom } T - \text{dom } T^*$  is a dense subspace of  $X$ . This supplies a simpler example for showing that the constraint qualification in the sum problem of maximal monotone operators can not be substantially weakened, see [14, Example 7.4]. Similarly, by Theorems 4.6 and 4.8,  $T_i^* = -T_i$ , we conclude that  $T_i + T_i^* = 0$  on  $\text{dom } T_i$ , a dense subset of  $L^2[0, 1]$ ; thus,  $T_i + T_i^*$  fails to be maximal monotone while both  $T_i, T_i^*$  are maximal monotone.

To study Fitzpatrick functions of sums of maximal monotone operators, we need:

**Lemma 4.13** *Let  $V$  be the Volterra integration operator (see Example 4.4). Then*

$$q_{V_+}^*(z) = \iota_{\text{span}\{e\}}(z) + \langle z, e \rangle^2, \quad \forall z \in X,$$

where  $e \equiv 1 \in L^2[0, 1]$ .

*Proof.* Let  $z \in X$ . By Example 4.4(iv) and Fact 2.5, we have

$$q_{V_+}^*(z) = \infty, \quad \text{if } z \notin \text{span}\{e\}.$$

Now suppose that  $z = le$  for some  $l \in \mathbb{R}$ . By Example 4.4(iv),

$$\begin{aligned} q_{V_+}^*(z) &= \sup_{x \in X} \{ \langle x, z \rangle - q_{V_+}(x) \} = \sup_{x \in X} \{ \langle x, le \rangle - \frac{1}{4} \langle x, e \rangle^2 \} \\ &= l^2 = \langle le, e \rangle^2 = \langle z, e \rangle^2. \end{aligned}$$

Hence  $q_{V_+}^*(z) = \iota_{\text{span}\{e\}}(z) + \langle z, e \rangle^2$ . ■

**Lemma 4.14** *Let  $T$  be defined as in Theorem 4.5. We have*

$$(35) \quad \begin{aligned} F_T(x, y^*) &= F_V(y^*, x) = \iota_{\text{span}\{e\}}(x + V^*y^*) + \frac{1}{2} \langle x + V^*y^*, e \rangle^2, \\ F_{T^*}(x, y^*) &= F_{V^*}(y^*, x) = \iota_{\text{span}\{e\}}(x + Vy^*) + \frac{1}{2} \langle x + Vy^*, e \rangle^2, \quad \forall (x, y^*) \in X \times X. \end{aligned}$$

*Proof.* Apply Fact 2.4, Fact 2.5 and Lemma 4.13. ■

**Remark 4.15** Theorem 4.16 below gives another example showing that  $F_{T+T^*} \neq F_T \square_2 F_{T^*}$  while  $T, T^*$  are maximal monotone, and  $\text{dom } T - \text{dom } T^*$  is a dense subspace in  $L^2[0, 1]$ . Moreover,  $\text{ran}(T_+ + (T^*)_+) = \{0\}$ . This again shows that the assumption that  $\text{dom } A - \text{dom } B$  is closed in Fact 2.8(ii) can not be weakened substantially, and that Fact 2.8(i) fails for discontinuous linear monotone operators.

**Theorem 4.16** *Let  $T$  be defined as in Theorem 4.5, and set*

$$H := \{x \in L^2[0, 1] : x \text{ is absolutely continuous, and } x' \in L^2[0, 1]\}.$$

*Then*

$$(36) \quad \begin{aligned} F_{T+T^*}(x, x^*) &= \iota_{X \times \{0\}}(x, x^*), \quad \forall (x, x^*) \in X \times X \\ F_T \square_2 F_{T^*}(x, x^*) &= \begin{cases} \frac{1}{2} [x(1)^2 + x(0)^2], & \text{if } (x, x^*) \in H \times \{0\}; \\ \infty, & \text{otherwise.} \end{cases} \end{aligned}$$

*Consequently,  $F_T \square_2 F_{T^*} \neq F_{T+T^*}$ .*

*Proof.* By Theorem 4.5(i) and Example 4.4(iii),

$$(37) \quad (T + T^*)y = 0, \forall y \in \text{dom } T \cap \text{dom } T^* = \{Vx \mid x \in e^\perp\},$$

where  $e \equiv 1 \in L^2[0, 1]$ . Let  $(x, x^*) \in X \times X$ . Using Theorem 4.5(i), we see that

$$(38) \quad F_{T+T^*}(x, x^*) = \sup_{y \in \text{dom } T \cap \text{dom } T^*} \langle x^*, y \rangle = \sup_{y \in X} \langle x^*, y \rangle = \iota_{\{0\}}(x^*) = \iota_{X \times \{0\}}(x, x^*).$$

By Fact 2.7, we have

$$(39) \quad (F_T \square_2 F_{T^*})(x, x^*) = \infty, \quad \forall x^* \neq 0.$$

When  $x^* = 0$ , by (35),

$$(40) \quad \begin{aligned} (F_T \square_2 F_{T^*})(x, 0) &= \inf_{y^* \in X} \{F_T(x, y^*) + F_{T^*}(x, -y^*)\} \\ &= \inf_{y^* \in X} \{\iota_{\text{span}\{e\}}(x + V^*y^*) + \frac{1}{2}\langle x + V^*y^*, e \rangle^2 + \iota_{\text{span}\{e\}}(x - Vy^*) + \frac{1}{2}\langle x - Vy^*, e \rangle^2\}. \end{aligned}$$

Observe that

$$\begin{aligned} x + V^*y^* \in \text{span}\{e\}, x - Vy^* \in \text{span}\{e\} \\ \Leftrightarrow x - Vy^* + Vy^* + V^*y^* \in \text{span}\{e\}, x - Vy^* \in \text{span}\{e\} \\ \Leftrightarrow x - Vy^* \in \text{span}\{e\}, \quad (\text{by Example 4.4(iv)}) \\ \Leftrightarrow x \in Vy^* + \text{span}\{e\} \Leftrightarrow x \text{ is absolutely continuous and } y^* = x'. \end{aligned}$$

Therefore,  $(F_T \square_2 F_{T^*})(x, 0) = \infty$  if  $x \notin H$ . For  $x \in H$ , using (40) and the fact that  $x - Vx' = x(0)e$  and  $x + V^*x' = x(1)e$ , we obtain

$$\begin{aligned} (F_T \square_2 F_{T^*})(x, 0) &= \frac{1}{2}\langle x + V^*x', e \rangle^2 + \frac{1}{2}\langle x - Vx', e \rangle^2 \\ &= \frac{1}{2}x(1)^2 + \frac{1}{2}x(0)^2 = \frac{1}{2}[x(1)^2 + x(0)^2]. \end{aligned}$$

Thus, (36) holds. Consequently,  $F_T \square_2 F_{T^*} \neq F_{T+T^*}$ . ■

Finally, we remark that the examples given in Sections 3 and 4 have important consequences on *decompositions of monotone operator*, namely Borwein-Wiersman decomposition and Asplund decomposition [7]. This will be addressed in the forthcoming paper [6].

## Acknowledgment

Heinz Bauschke was partially supported by the Natural Sciences and Engineering Research Council of Canada and by the Canada Research Chair Program. Xianfu Wang was partially supported by the Natural Sciences and Engineering Research Council of Canada.

## References

- [1] H.H. Bauschke and J.M. Borwein, “Maximal monotonicity of dense type, local maximal monotonicity, and monotonicity of the conjugate are all the same for continuous linear operators”, *Pacific Journal of Mathematics*, vol. 189, pp. 1–20, 1999.
- [2] H.H. Bauschke, J.M. Borwein, and X. Wang, “Fitzpatrick functions and continuous linear monotone operators”, *SIAM Journal on Optimization*, vol. 18, pp. 789–809, 2007.
- [3] H.H. Bauschke, D.A. McLaren, and H.S. Sendov, “Fitzpatrick functions: inequalities, examples and remarks on a problem by S. Fitzpatrick”, *Journal of Convex Analysis*, vol. 13, pp. 499–523, 2006.
- [4] H.H. Bauschke, X. Wang, and L. Yao, “Autoconjugate representers for linear monotone operators”, *Mathematical Programming (Series B)*, vol. 123, pp. 5–24, 2010.
- [5] H.H. Bauschke, X. Wang, and L. Yao, “Monotone linear relations: maximality and Fitzpatrick functions”, *Journal of Convex Analysis*, vol. 16, pp. 673–686, 2009.
- [6] H.H. Bauschke, X. Wang, and L. Yao, “On Borwein-Wiersma Decompositions of monotone linear relations”, <http://arxiv.org/abs/0912.2772>, December 2009.
- [7] J.M. Borwein and H. Wiersma, “Asplund decomposition of monotone operators”, *SIAM Journal on Optimization*, 18, pp. 946–960, 2007.
- [8] H. Brézis, *Opérateurs Maximaux Monotones et Semi-Groupes de Contractions dans les Espaces de Hilbert*, North-Holland Publishing, Amsterdam, 1973.
- [9] H. Brézis and F.E. Browder, “Linear maximal monotone operators and singular nonlinear integral equations of Hammerstein type”, in *Nonlinear analysis (collection of papers in honor of Erich H. Rothe)*, Academic Press, pp. 31–42, 1978.
- [10] R. Cross, *Multivalued Linear Operators*, Marcel Dekker, Inc, New York, 1998.
- [11] F. Deutsch, *Best Approximation in Inner Product Spaces*, Springer-Verlag, New York, 2001.
- [12] S. Fitzpatrick, “Representing monotone operators by convex functions”, in *Workshop/Miniconference on Functional Analysis and Optimization (Canberra 1988)*, Proceedings of the Centre for Mathematical Analysis, Australian National University vol. 20, Canberra, Australia, pp. 59–65, 1988.
- [13] P.R. Halmos, *A Hilbert Space Problem Book*, Van Nostrand Reinbold, 1967.
- [14] R.R. Phelps and S. Simons, “Unbounded linear monotone operators on nonreflexive Banach spaces”, *Journal of Convex Analysis*, vol. 5, pp. 303–328, 1998.
- [15] J.-P. Penot, “The relevance of convex analysis for the study of monotonicity”, *Nonlinear Analysis*, vol. 58, pp. 855–871, 2004.



- [16] R.T. Rockafellar, “On the maximality of sums of nonlinear monotone operators”, *Transactions of the American Mathematical Society*, 149 pp. 75–88, 1970.
- [17] W. Rudin, *Functional Analysis*, McGraw-Hill, 2nd ed., 1991.
- [18] S. Simons, *From Hahn-Banach to Monotonicity*, Springer-Verlag, 2008.
- [19] S. Simons and C. Zălinescu, “Fenchel duality, Fitzpatrick functions and maximal monotonicity”, *Journal of Nonlinear and Convex Analysis* vol. 6, pp. 1–22, 2005.
- [20] K.R. Stromberg, *An Introduction to Classical Real Analysis*, Wadsworth, 1981.
- [21] B.F. Svaiter, “Non-enlargeable operators and self-cancelling operators”, *Journal of Convex Analysis*, vol. 17, pp. 309–320, 2010.
- [22] M.D. Voisei, “The sum theorem for linear maximal monotone operators”, *Mathematical Sciences Research Journal*, vol. 10, pp. 83–85, 2006.
- [23] M.D. Voisei and C. Zălinescu, “Linear monotone subspaces of locally convex spaces”, *Set-Valued and Variational Analysis*, vol. 18, pp. 29–55, 2010.
- [24] E. Zeidler, *Nonlinear Functional Analysis and its Application, Vol II/A Linear Monotone Operators*, Springer-Verlag, New York-Berlin-Heidelberg, 1990.
- [25] E. Zeidler, *Nonlinear Functional Analysis and its Application, Vol II/B Nonlinear Monotone Operators*, Springer-Verlag, New York-Berlin-Heidelberg, 1990.